

Tibial Plateau Leveling Osteotomy

The cranial cruciate ligament (CCL) in the dog is one of the major stabilizers of the stifle (knee). It serves the same function as the anterior cruciate ligament in people, preventing internal rotation of the stifle, forward displacement of the tibia (the shin bone), and hyperextension.

The cranial cruciate ligament (CCL) in the dog is one of the major stabilizers of the stifle (knee). It serves the same function as the anterior cruciate ligament in people, preventing internal rotation of the stifle, forward displacement of the tibia (the shin bone), and hyperextension. It can tear during a sudden trauma, or more commonly it slowly deteriorates over weeks or months. Following a CCL tear in dogs, it is possible for the other stifle to develop a CCL tear as well within the near future. Following a CCL tear, the limp is usually severe. While some improvement can occur with time, surgery is the best method to resolve the lameness and to minimize future osteoarthritis.

There have been numerous surgical techniques developed over the past 30 years. The most recently developed and most promising techniques for restoring normal limb use involve tibial osteotomy (bone cutting). The osteotomy techniques alter the weightbearing forces acting at the stifle. Instead of repairing the CCL, these techniques focus on eliminating the need for the CCL. Tibial plateau leveling osteotomy (TPLO) and tibial tuberosity advancement (TTA) and are the two most commonly performed tibial osteotomy techniques.

TPLO was developed in the early 1990s, and in recent years it has become an increasingly popular technique. Initially developed for large breed dogs, TPLO is now performed in dogs (and some cats) of all sizes. With this surgery, the cranial cruciate ligament is not repaired. Instead, the tibial plateau (the part of the tibia just below the stifle) is repositioned to counteract the instability which occurs following the cruciate ligament injury. A specially designed bone plate with screws is used to hold the bone in the new position until healing occurs. While the surgery is more involved than previous repair techniques, patient comfort is very good during recovery. Also, most dogs will become weight bearing on the leg sooner after TPLO compared to older methods of repair.

Vienna

The Hope Center
140 Park St, SE
Vienna, VA 22180
p 703.242.6000
f 703.242.8358

Leesburg

The LifeCentre
165 Fort Evans Rd, NE
Leesburg, VA 20176
p 703.771.2100
f 703.777.9968

Winchester

Valley Veterinary Emergency
and Referral Clinic
164-4 Garber Lane
Winchester, VA 22602
p 540.450.0177
f 703.777.9968

Woodbridge

Woodbridge Animal Hospital
2703 Caton Hill Road
Woodbridge, VA 22191
p 703.242.6000
f 703.242.8358

Strict rest is required following surgery to allow for proper healing of the tibia. Initially, there should be no running, jumping, or playing. After the first month of strict rest, a gradual introduction of controlled activity is performed until 3 months following surgery. Physical therapy is also performed to improve the speed of healing. In the weeks following surgery, x-rays are taken to assess healing of the tibia.



Vienna

The Hope Center
140 Park St, SE
Vienna, VA 22180
p 703.242.6000
f 703.242.8358

Leesburg

The LifeCentre
165 Fort Evans Rd, NE
Leesburg, VA 20176
p 703.771.2100
f 703.777.9968

Winchester

Valley Veterinary Emergency
and Referral Clinic
164-4 Garber Lane
Winchester, VA 22602
p 540.450.0177
f 703.777.9968

Woodbridge

Woodbridge Animal Hospital
2703 Caton Hill Road
Woodbridge, VA 22191
p 703.242.6000
f 703.242.8358

